
Bedside Medicine Without Tears 2nd Edition

bedside ultrasound for detection of deep vein thrombosis ... - two-point compression method to detect dvt israeli journal of emergency medicine - vol. 6, no. 3 sept. 2006 - בתכ תעה ילארשיה האופרל הפוחד 36 **culture shock — patient as icon, icon as patient** - n engl j med 359;26 nejm december 25, 2008 perspective 2749 bulbar palsy — which today are uncommonly recognized. when i stroked a patient's palm and **technique for safe placement of a dobhoff tube without a ...** - 66 | the medicine forum 5 technique for safe placement of a dobhoff tube without a cortrak® machine jon chao, md, jennifer alloo hong, md a 58 year old female with a history of end stage renal **obstetric emergencies - american association of physician ...** - 44 american journal of clinical medicine® • spring 2009 • volume six, number two obstetric emergencies tried replacing the cord into the uterus, it is not successful. **espen guidelines on enteral nutrition: liver disease** - clinical nutrition (2006) 25, 285-294 espen guidelines espen guidelines on enteral nutrition: liver disease\$ m. plautha,, e. cabre´b, o. riggioc, m. assis-camilod, **ethical considerations in emergency care - isrjem** - emergency medicine ethics behind modern developments, the bioethical construct allows a greater flexibility in decision making. this is a crucial factor in the crucible of the emergency room, **glycemic control in critical care - ihi** - details elements develop and implement a method of measuring and tracking glucose levels and displaying them for front-line staff. • determine which glucose values to use: bedside glucometer values, laboratory values, or both. **the 4a test: screening - guy's and st thomas** - title: short orientation-memory-concentration (omc) test of cognitive function author: maclulich created date: 9/9/2011 7:30:57 am **simplified criteria for the diagnosis of autoimmune hepatitis** - autoimmune, cholestatic and biliary disease simplified criteria for the diagnosis of autoimmune hepatitis elkem.hennes,1 mikiozeniya,2 albert.jaja,3 albertpares,´ 4 georgen.dalekos,5 edwardlawitt,6 **the short confusion assessment method (short cam ...** - 2 background delirium (acute confusional state) is a common, serious, and potentially preventable source of morbidity and mortality for older hospitalized patients. **cleaning can be programmatically improved** - cleaning can be programmatically improved 40 50 60 70 80 % of objects cleaned pre intervention. post intervention. p =